

Supplementary Table 1. Diagnosis of congenital heart disease according to the presence of pulmonary hypertension

	Pulmonary hypertension (n= 15)	No pulmonary hypertension (n = 61)	Total (n = 76)	P-value
Diagnosis				0.064
Ventricular septal defect	11 (73.3)	19 (31.1)	30(39.5)	
Tetralogy of Fallot	0 (0)	22 (36.1)	22 (28.9)	
Atrioventricular canal defect	2 (13.3)	2 (3.3)	4 (5.3)	
Coarctation of aorta	0 (0.0)	3 (4.9)	3 (3.9)	
Coarctation of aorta with ventricular septal defect	0 (0)	2 (3.3)	2 (2.6)	
Atrial septal defect	1 (6.7)	2 (3.3)	3 (3.9)	
Double outlet right ventricle	0 (0.0)	3 (4.9)	3 (3.9)	
Transposition of great arteries with intact ventricular septum	0 (0.0)	2 (3.3)	2 (2.6)	
Left ventricular out tract obstruction	0 (0.0)	2 (3.3)	2 (2.6)	
Tricuspid regurgitation	0 (0.0)	1 (1.6)	1 (1.3)	
Ebstein's anomaly	0 (0.0)	1 (1.6)	1 (1.3)	
Right ventricular out tract obstruction	0 (0.0)	1 (1.6)	1 (1.3)	
Partial anomalous pulmonary venous return	1 (6.7)	0 (0.0)	1 (1.3)	
Interrupted aortic arch with ventricular septal defect	0 (0.0)	1 (1.6)	1 (1.3)	