



Corrigendum

Korean J Anesthesiol 2021;74(2):188
<https://doi.org/10.4097/kja.19305.e1>
pISSN 2005-6419 • eISSN 2005-7563


Fluid responsiveness in the pediatric population

Ji-Hyun Lee, Eun-Hee Kim, Young-Eun Jang, Hee-Soo Kim, Jin-Tae Kim

Department of Anesthesiology and Pain Medicine, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, Korea

Korean J Anesthesiol 2019 October 72(5): 429-440
<https://doi.org/10.4097/kja.19305>

The article by Lee et al. entitled, “Fluid responsiveness in the pediatric population” (Korean J Anesthesiol 2019 Oct; 72(5): 429-440) should add the following explanation in Fig. 2.



The unit of the suggested equations for PPV and SPV is %.

Although SPV (mmHg) is traditionally calculated as $SP_{max} - SP_{min}$ or $\Delta Up + \Delta Down$, SPV(%) has been used for pediatric patients.

© The Korean Society of Anesthesiologists, 2021

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.