

Supplementary Table 1. Questionnaire about the experience at 24 hours postoperatively stratified by group.

	CCB20 (n=30)	CCB40 (n=30)	<i>p</i> -value
Least pain score ^a	2.0 (1.0, 4.0)	3.0 (2.0, 4.0)	0.327
Worst pain score ^a	7.5 (5.0, 9.0)	7.0 (3.0, 8.0)	0.127
Proportion of time with severe pain ^b	5.0 (3.0,7.0)	4.0 (2.0,7.0)	0.213
Sleep disturbance due to pain ^a	5.5 (3.0, 9.0)	4.5 (2.0, 7.0)	0.288
Nausea ^a	0.0 (0.0, 2.0)	0.0 (0.0, 1.0)	0.979
Vomiting ^a	0.0 (0.0, 0.0)	0.0 (0.0, 0.0)	0.211
Drowsiness ^a	1.5 (0.0, 3.0)	1.5 (0.0, 5.0)	0.779
Pruritus ^a	0.0 (0.0, 0.0)	0.0 (0.0, 0.0)	0.106
Dizziness ^a	0.0 (0.0, 2.0)	0.0 (0.0, 1.0)	0.748
Global Satisfaction ^c	7.0 (5.0, 9.0)	8.0 (6.0, 9.0)	0.407

Values are presented as median (1Q, 3Q). CCB20, costoclavicular block (CCB) with 20 mL of 0.75% ropivacaine; CCB40, CCB with 40 mL of 0.375% ropivacaine. All values are numeric rating scales (0 to 10). ^a 0, best; 10, worst. ^b How often do you feel severe pain during postoperative 24 hours? 0, least; 10, most. ^c 0, worst; 10, best.