



Relationship between first-twitch depression and train-of-four ratio during sugammadex reversal of rocuronium-induced neuromuscular blockade

Kyo Sang Kim, You Na Oh, Tae Yeon Kim, Song Yee Oh, and Yeong Hun Sin

Department of Anesthesiology and Pain Medicine, Hanyang University College of Medicine, Seoul, Korea

Background: The primary outcome of sugammadex reversal for rocuronium-induced neuromuscular block (NMB) is a train-of-four ratio (TOFR) of 0.9, not first twitch (T_1) height. We investigated whether the recovery of TOFR or T_1 differs based on the reversal of NMB with neostigmine or sugammadex.

Methods: The acceleromyographic responses from 0.6 mg/kg of rocuronium were monitored supramaximally in 80 patients after induction of anesthesia. The TOFR and T_1 height were recorded, and saved in a personal computer using TOF-Watch SX Monitor software in all patients. Patients were randomly assigned to 2 groups to receive either neostigmine 50 μ g/kg with glycopyrrolate 10 μ g/kg (neostigmine group, $n = 40$) or sugammadex 2.0 mg/kg (sugammadex group, $n = 40$). The primary objective was to determine the difference of recovery time between TOFR to 0.9 and T_1 to 0.9 after sugammadex or neostigmine administration during moderate rocuronium-induced NMB.

Results: The recovery pattern of the TOFR 2 min after sugammadex administration was 1.0 or more, but that of T_1 was less than 90% (T_1 / control value) up to 6 min after drug was injected. The recovery pattern of TOFR and T_1 was similar during the 20 min after reversal with neostigmine.

Conclusions: If you have not performed the T_1 monitoring, both TOFR and T_1 should be considered to confirm suitable recovery during the 6 min after reversal with sugammadex during rocuronium-induced moderate NMB.

Key Words: Neostigmine, Neuromuscular monitoring, Rocuronium, Sugammadex.

Introduction

Exact monitoring of muscle relaxation is essential for proper use of muscle relaxants and to prevent post-operative residual paralysis [1]. Train-of-four (TOF) monitoring and single twitch (ST) stimulation of peripheral nerves is often used to monitor the degree of neuromuscular block (NMB) when anesthetic techniques include use of neuromuscular blocking agents (NMBAs). Using ST should establish a control value before the administration of NMBAs. ST and first twitch of TOF (T_1) forces do not differ when employing a stimulus interval of 10 seconds (s) or more [2]. The T_1 response represents the effects of NMBAs in the postsynaptic membrane, while the TOF ratio

Corresponding author: Kyo Sang Kim, M.D., Ph.D.
Department of Anesthesiology and Pain Medicine, Hanyang University
College of Medicine, 222, Wangsimni-ro, Seongdong-gu, Seoul 04763,
Korea
Tel: 82-2-2290-8680, Fax: 82-2-2299-0742
E-mail: kimks@hanyang.ac.kr
ORCID: <http://orcid.org/0000-0002-2186-3484>

Received: January 13, 2016.
Revised: 1st, February 13, 2016; 2nd, March 8, 2016.
Accepted: March 14, 2016.

Korean J Anesthesiol 2016 June 69(3): 239-243
<http://dx.doi.org/10.4097/kjae.2016.69.3.239>

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © the Korean Society of Anesthesiologists, 2016

Online access in <http://ekja.org>

(T_4/T_1 , TOFR) shows the effect on the pre-synaptic membrane of the neuromuscular junction [3].

During spontaneous or pharmacologically-induced offset of NMB, a TOFR of 0.9 has been reported to exclude clinically important residual NMB [3]. The T_1 usually recovers at a rate similar to or faster than the TOFR after the reversal of competitive NMBAs by neostigmine. After reversal with the optimum dose of sugammadex during rocuronium-induced NMB, the time to 0.9 of the TOFR is much faster than that to 90% of T_1 (T_1 / control value). A TOFR of 0.9 cannot assure adequate recovery of neuromuscular function in several minutes after sugammadex administration, if the T_1 was not recovered completely [4,5]. However, this previous report simply described the time of a T_1 and TOFR to 0.9, not saved in a personal computer using TOF-Watch SX Monitor software. So the detailed relationship between a TOFR and T_1 should be analyzed following the time after sugammadex administration.

We analyzed the relationship of T_1 recovery and TOFR after administration of sugammadex 2.0 mg/kg or neostigmine 50 µg/kg during rocuronium-induced moderate NMB.

Materials and Methods

After approval of the protocol by the Hospital Ethics Committee, 80 adult patients were enrolled in this study and all granted written informed consent. This randomized, parallel-group, blinded study was conducted in accordance with principles of Good Clinical Research Practice [3]. All patients of both sexes, American Society of Anesthesiologists physical status I–II, aged between 20–64 years, received elective surgery under

general anesthesia with rocuronium for intubation and maintenance. Using a computer-generated program, patients were randomly assigned to either the neostigmine group (n = 40, neostigmine 50 µg/kg with glycopyrrolate 10 µg/kg after operation) or the sugammadex group (n = 40, sugammadex 2.0 mg/kg after operation) (Fig. 1). Exclusion criteria were as follows: predicted difficult intubation, previous known neuromuscular diseases that may affect NMB, allergy to any drug used in general anesthesia, history of serious liver or kidney disease, use of drugs that might interact with NMBAs, pregnancy, or obesity (defined as a body mass index [BMI] ≥ 30 kg/m²).

In the operating room, the patients received a standard monitoring of pulse oximetry, capnography, electrocardiography and noninvasive blood pressure measurement. Depth of hypnosis was evaluated for all patients using a bispectral index (BIS) XP monitor (Model A 2000, Aspect Medical Systems, Newton, MA, USA). Anesthesia was induced without premedication, using 2.0–2.5 mg/kg propofol, 0.5 µg/kg/min remifentanyl, and oxygen, and was maintained with 1.0–1.5 MAC of sevoflurane in 50% air in oxygen and 0.1–0.3 µg/kg/min remifentanyl to maintain a BIS monitor reading between 40 and 50 throughout surgery. During maintenance of anesthesia, end-tidal PCO₂ was kept between 30 and 35 mmHg.

NMB was monitored using a TOF-Watch SX[®] (Merck Sharp & Dohme Corp., Glostrup, Denmark) at the adductor pollicis muscle. Two surface electrodes (Cleartrode[™], ConMed[®], Utica, NY, USA) were placed on cleaned skin overlying the ulnar nerve at the wrist. Stabilization was achieved with a 5 s, 50 Hz tetanic stimulation. Supramaximal stimuli were applied after automatic calibration (CAL 2 mode) of acceleromyography after an initial




Fig. 1. Patient flow through the study. T₂: second twitch reappeared in TOF stimulation.

tetanic stimulus [3]. If first TFR and T_1 were $< 95\%$ or $> 105\%$, we performed the recalibration after the change of the location of a piezoelectric transducer or fixed wrist. This experiment was begun after confirming a TFR and T_1 of 95%–105%. The TOF mode of supramaximal stimulation (0.2 ms duration, frequency 2 Hz, 2 s duration) was applied at 15 s intervals, which lasted until the end of anesthesia. After stabilization of control responses, rocuronium was administered at 0.6 mg/kg before tracheal intubation, then subsequently at 0.2 mg/kg when the second twitch (T_2) reappeared in TOF stimulation.

The primary objective of this study was to determine recovery time and response after sugammadex or neostigmine administration of the T_1 and TOFR to 0.9 during rocuronium-induced moderate NMB. After the surgical dressing, each patient received sugammadex 2.0 mg/kg or a single dose of neostigmine 50 $\mu\text{g}/\text{kg}$ plus glycopyrrolate 10 $\mu\text{g}/\text{kg}$, respectively, when T_2 reappeared in TOF stimulation. All neuromuscular monitoring data were saved in a personal computer using TOF-Watch SX Monitor software. Full recovery from NMB was conducted during the administration of sevoflurane and remifentanyl. Extubation was carried out after consciousness and regular respiration were confirmed. Skin temperature of the hand was measured and kept above 32°C. Central temperature was continuously monitored at the lower esophagus and kept above 35.5°C.

Statistical analysis

Statistical analysis was performed with SPSS 17.0 for Windows (SPSS Inc., Chicago, IL, USA). Sample size was determined based on the difference of recovery time (mean [SD] recovery times of 4.4 [7.9] min) between TOFR to 0.9 and T_1 to 0.9 after sugammadex 4 mg/kg during deep rocuronium-induced NMB [5]. We considered time reduction by more than 40%, to 2.6 min (SD 3.2), to be clinically significant. Obtaining statistically significant results with $\alpha = 0.05$ and a power of 0.8 required 37 patients. Projecting 10% dropout, we enrolled 40 patients in each group. Characteristics of patients were compared using the Chi square test between the two groups, and T_1 height and TOFR

were compared using an unpaired t-test. Repeated measures analyses of variance (ANOVA) were conducted for identifying the neuromuscular recovery. Data are expressed as the mean \pm SD, range, or number. Differences were considered to be significant at $P < 0.05$.

Results

All enrolled patients completed the study, and neuromuscular monitoring was successfully conducted. The two groups were well balanced in their demographic profile, end-tidal concentration of sevoflurane and anesthetic time (Table 1). The T_1 recovery and TOFR was 87.8% \pm 17.5% and 90.1% \pm 12.1% at 15 min after neostigmine 50 $\mu\text{g}/\text{kg}$, respectively ($P = 0.496$). The recovery pattern of T_1 and TOFR was similar during the 20 min after neostigmine administration (Fig. 2). The T_1 recovery and TOFR was 90.4% \pm 13.5% and 104.9% \pm 6.7% at 6 min after sugammadex 2 mg/kg, respectively ($P < 0.001$). The recovery pattern of TOFR after 2 min was 100% or more, but that of T_1 was less than 90% up to 6 min after sugammadex administration (Fig.




Fig. 2. Progress of the first twitch and the train-of-four ratio after administration of neostigmine during rocuronium-induced neuromuscular block. Neostigmine is administered at time 0 when the second twitch reappears in TOF stimulation.

Table 1. Demographic Data and Anesthetic Drug Parameters

	Neostigmine	Sugammadex	P
N	40	40	
Gender (M/F)	22/18	24/16	0.657
Age (yr)	42.2 \pm 12.2 (21–64)	40.5 \pm 13.1 (20–62)	0.550
Weight (kg)	64.0 \pm 10.5	64.8 \pm 12.7	0.760
Height (cm)	164.3 \pm 7.6	165.4 \pm 9.3	0.564
Sevoflurane at antagonism (vol%, end-tidal)	0.4 \pm 0.3	0.5 \pm 0.3	0.140
Anesthetic time (min)	48.6 \pm 20.3	52.9 \pm 25.2	0.403

Values are mean \pm SD, range, or number.




Fig. 3. Progress of the first twitch and the train-of-four ratio after administration of sugammadex during rocuronium-induced neuromuscular block. Sugammadex is administered at time 0 when the second twitch reappears in TOF stimulation. *P < 0.001 versus first twitch. †P < 0.01 versus first twitch.

3). Of particular note, the neuromuscular recovery of T₁ at 120 s following administration of sugammadex was only 80.2% ± 16.7%, but the TOFR at the same time was 104.8% ± 8.4% (P < 0.001) (Table 2). The T₁ recovery was 12.3% ± 6.1% and 15.2% ± 7.8% in the neostigmine and sugammadex groups, respectively, at time 0 when the second twitch was observed in TOF stimulation (P = 0.257) (Figs. 2 and 3).

Discussion

We found that the recovery pattern of the TOFR after 2 min was 1.0 or more, but that of T₁ was less than 90% up to 6 min after sugammadex administration. In particular, the neuromuscular recovery of T₁ was only 80% compared with a TOFR of 1.0 at 120 s. The recovery pattern of T₁ and TOFR was similar during the 20 min after neostigmine administration.

Postoperative residual paralysis is frequent, dangerous, and difficult to recognize clinically. An impaired ventilatory response to hypoxemia, visual impairment and the possibility of increased risk of aspiration occurs when the TOFR is < 0.9. Patients are likely to feel an unpleasant feeling of residual paralysis in a short time [6]. Even with recovery to a TOFR of more than 0.9, the incidence of pharyngeal dysfunction is 13% [7]. It is recommended that a TOFR of more than 0.9 should be confirmed before extubation using acceleromyographic monitoring.

The relationship between the absolute value of T₁ height and the TOFR is constant and reliable as a method of monitoring the effects of NMBA and their antagonists [8]. When T₁ height has returned to a control value, the recovery pattern of TOF stimulation frequently fades markedly (indicating residual receptor occupancy). T₁ recovery to 90% is similar to a TOFR of 0.7,

Table 2. Neuromuscular Recovery (%) from Administration of Sugammadex in First Twitch and Train-of-four Ratio during 120 Seconds (s)

	First twitch	Train-of-four ratio	P
Injection	15.2 ± 7.8	0	< 0.001
15 s	18.6 ± 7.2	9.0 ± 16.3	0.001
30 s	35.5 ± 12.7	47.6 ± 29.5	0.02
45 s	53.1 ± 11.4	70.2 ± 24.0	< 0.001
60 s	65.8 ± 12.8	88.2 ± 14.0	< 0.001
75 s	70.7 ± 14.1	98.4 ± 12.1	< 0.001
90 s	75.2 ± 15.6	104.6 ± 11.1	< 0.001
105 s	77.8 ± 15.6	104.5 ± 10.3	< 0.001
120 s	80.2 ± 16.7	104.8 ± 8.4	< 0.001

representing blockage of 80% of neuromuscular receptors after anticholinesterase or spontaneous recovery [9,10]. T₁ height recovers to 75% of control at a TOFR of 0.25, and returns to 90% of control when the TOFR exceeds 0.9 in electromyographic records [8]. However, in acceleromyographic monitoring, T₁ returns to less than 75% of control on average when the TOFR is 0.7. When the TOFR returns to 0.9, T₁ is still less than 90% of control [11]. We confirmed a similar relationship between T₁ and TOFR after reversal of neostigmine in acceleromyographic recordings (Fig. 2).

Results differ between mechanomyography and acceleromyography at approximately 50% twitch height maximally, with acceleromyography frequently overestimating the extent of the block [12]. The overestimated data from acceleromyographic monitoring should be normalized by baseline values to reliably detect residual paralysis [13]. Data normalization was not necessary in the present study, because TOFR and T₁ recovery were recorded in a TOF response in each patient.

TOFR shows significantly more rapid recovery than T₁ height after an optimum dose of sugammadex [4,5]. If the administered dose of sugammadex is small, the recovery of the TOFR to 0.9 appears significantly slower than recovery of T₁ to 90% [5]. A TOFR of 0.9 may not guarantee an appropriate recovery of neuromuscular function if T₁ is still less than 90% after sugammadex administration. We found that after sugammadex 2 mg/kg during rocuronium-induced moderate NMB, the T₁ height recovered to 90% by 6 min after administration, although TOFR was already recovered to 0.9. Furthermore, since the recovered T₁ was less than 80% until 2 min after administration, a TOFR of 1.0 may not be sufficient for the quick and appropriate recovery of neuromuscular relaxation (Table 2). If the inhalation and intravenous anesthetics are early stopped before the administration of sugammadex to reduce the recovery time, the patient may be in the awakening before complete recovery from muscle relaxation. At that time, the patient is likely to experience an unpleasant feeling due to residual paralysis, which causes a sud-

den and harsh response in the recovery process. We suspect this symptom is happened due to still depression of a T_1 compared with full recovery of a TOFR. Maintenance of sevoflurane until administration of sugammadex allows the time for washing out of the inhalation anesthetic to awaken.

The reason for the difference between TOFR and first twitch after reversal with sugammadex is unclear, but we suspect that sugammadex results in a drastic reduction of the effective site concentration of the unbound relaxant due to formation of a stable complex with rocuronium in the plasma. Through a gradient of rocuronium molecules between the plasma and neuromuscular junction, NMBAs diffuse away from the nicotinic receptor, providing rapid recovery from NMB and prompt recovery of TOFR. However, in contrast to the TOFR, the process does not promote reestablishment of T_1 with the same proportion and speed, i.e., muscle receptors are still blocked by rocuronium in the short term [14].

The previous reports [4,5] compared simply the time of a T_1 and TOFR to 0.9, not described the specific pattern of recovery percent after sugammadex during rocuronium-induced deep NMB. In the present study, however, all neuromuscular moni-

toring data were saved in a personal computer using TOF-Watch SX Monitor software. So the detailed relationship between a TOFR and T_1 can be analyzed following the time after sugammadex administration.

This study contains some limitations. The present results are unique to patients undergoing moderate rocuronium-induced NMB, and may not be generalizable to patients undergoing deep or pronounced rocuronium-induced NMB. The current results are difficult to actually verify in patients during recovery from anesthesia, because it is not possible to distinguish the impact of anesthetics and opioids from those of NMBAs and reversal agents. Maintenance of sevoflurane delays the recovery from rocuronium compared with intravenous anesthesia. Even though this trial has been implemented in the same environment in each group, sevoflurane may have affected recovery from rocuronium.

In conclusion, both TOFR and T_1 should be considered when confirming adequate recovery in the 6 min after reversal with sugammadex administration during rocuronium-induced moderate NMB, if you have not conducted a separate T_1 monitoring.

References

1. Murphy GS, Szokol JW, Marymont JH, Greenberg SB, Avram MJ, Vender JS, et al. Intraoperative acceleromyographic monitoring reduces the risk of residual neuromuscular blockade and adverse respiratory events in the postanesthesia care unit. *Anesthesiology* 2008; 109: 389-98.
2. van Santen G, Fidler V, Houwertjes MC, Top WM, Wierda JM. The effect of single twitch and train-of-four stimulation on twitch forces during stable neuromuscular block. *J Clin Monit Comput* 2000; 16: 529-33.
3. Fuchs-Buder T, Claudius C, Skovgaard LT, Eriksson LI, Mirakhur RK, Viby-Mogensen J. Good clinical research practice in pharmacodynamic studies of neuromuscular blocking agents II: the Stockholm revision. *Acta Anaesthesiol Scand* 2007; 51: 789-808.
4. Suzuki T. A train-of-four ratio of 0.9 may not certify adequate recovery after sugammadex. *Acta Anaesthesiol Scand* 2011; 55: 368-9.
5. Staals LM, Driessen JJ, Van Egmond J, De Boer HD, Klimek M, Flockton EA, et al. Train-of-four ratio recovery often precedes twitch recovery when neuromuscular block is reversed by sugammadex. *Acta Anaesthesiol Scand* 2011; 55: 700-7.
6. Kopman AF, Yee PS, Neuman GG. Relationship of the train-of-four fade ratio to clinical signs and symptoms of residual paralysis in awake volunteers. *Anesthesiology* 1997; 86: 765-71.
7. Sundman E, Witt H, Olsson R, Ekberg O, Kuylenstierna R, Eriksson LI. The incidence and mechanisms of pharyngeal and upper esophageal dysfunction in partially paralyzed humans: pharyngeal videoradiography and simultaneous manometry after atracurium. *Anesthesiology* 2000; 92: 977-84.
8. Power SJ, Jones RM. Relationship between single twitch depression and train-of-four fade: influence of relaxant dose during onset and spontaneous offset of neuromuscular blockade. *Anesth Analg* 1987; 66: 633-6.
9. Waud BE, Waud DR. The relation between the response to "train-of-four" stimulation and receptor occlusion during competitive neuromuscular block. *Anesthesiology* 1972; 37: 413-6.
10. Lee CM. Train-of-4 quantitation of competitive neuromuscular block. *Anesth Analg* 1975; 54: 649-53.
11. Kopman AF, Klewicka MM, Neuman GG. The relationship between acceleromyographic train-of-four fade and single twitch depression. *Anesthesiology* 2002; 96: 583-7.
12. Harper NJ, Martlew R, Strang T, Wallace M. Monitoring neuromuscular block by acceleromyography: comparison of the Mini-Accelograph with the Myograph 2000. *Br J Anaesth* 1994; 72: 411-4.
13. Suzuki T, Fukano N, Kitajima O, Saeki S, Ogawa S. Normalization of acceleromyographic train-of-four ratio by baseline value for detecting residual neuromuscular block. *Br J Anaesth* 2006; 96: 44-7.
14. Sorgenfrei IF, Norrild K, Larsen PB, Stensballe J, Ostergaard D, Prins ME, et al. Reversal of rocuronium-induced neuromuscular block by the selective relaxant binding agent sugammadex: a dose-finding and safety study. *Anesthesiology* 2006; 104: 667-74.